

Manuals+

[Q & A](#) | [Deep Search](#) | [Upload](#)

manuals.plus /

› [Mucinex](#) /

› [Mucinex Fast-Max Severe Congestion & Cough Medicine User Manual](#)

Mucinex 6380000000

Mucinex Fast-Max Severe Congestion & Cough Medicine User Manual

Model: 6380000000

1. PRODUCT OVERVIEW

Mucinex Fast-Max Severe Congestion & Cough is an over-the-counter medication designed to provide maximum strength relief for severe cold and flu symptoms. This liquid formula targets cough, chest congestion, and nasal congestion, helping to thin and loosen mucus for easier breathing.

Key Benefits:

- Controls cough.
- Relieves nasal and chest congestion.
- Thins and loosens mucus to help clear lungs.

MAXIMUM STRENGTH SEVERE CONGESTION & COUGH RELIEF



COUGH



**NASAL
CONGESTION**



**CHEST
CONGESTION**

Image 1: Front view of the Mucinex Fast-Max Severe Congestion & Cough liquid medicine bottle.

2. IMPORTANT SAFETY INFORMATION

Warnings:

- Do not use for children under 12 years of age.
- Do not use if you are currently taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. Consult a doctor or pharmacist before use if unsure.

Consult a doctor before use if you have:

- Heart disease
- High blood pressure
- Thyroid disease
- Diabetes
- Trouble urinating due to an enlarged prostate gland

- Persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- Cough that occurs with too much phlegm (mucus)

When using this product:

- Do not use more than directed.

Stop use and ask a doctor if:

- Nervousness, dizziness, or sleeplessness occur.
- Symptoms do not get better within 7 days or occur with fever.
- Cough returns, or occurs with rash or persistent headache. These could indicate a serious condition.

Pregnancy/Breast-feeding: If pregnant or breast-feeding, consult a health professional before use.

Keep out of reach of children. In case of overdose, seek medical help or contact a Poison Control Center immediately.

Tamper Evident: Do not use if the neckband on the bottle cap is broken or missing.

MAXIMUM STRENGTH

9

DOSES

<p>Drug Facts</p> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 50%; text-align: left; border-bottom: 1px solid black;">Active ingredients (in each 20 mL)</th> <th style="width: 50%; text-align: left; border-bottom: 1px solid black;">Purposes</th> </tr> <tr> <td style="font-size: 0.8em;">Dextromethorphan HBr 20 mg...Cough suppressant Guaifenesin 400 mg.....Expectorant Phenylephrine HCl 10 mg.....Nasal decongestant</td> <td></td> </tr> </table> <p>Uses</p> <ul style="list-style-type: none"> ■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive ■ temporarily relieves: <ul style="list-style-type: none"> ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants ■ the intensity of coughing ■ the impulse to cough to help you get to sleep ■ nasal congestion due to a cold <p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ▶ 	Active ingredients (in each 20 mL)	Purposes	Dextromethorphan HBr 20 mg...Cough suppressant Guaifenesin 400 mg.....Expectorant Phenylephrine HCl 10 mg.....Nasal decongestant		<p>Drug Facts (continued)</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus) <p>When using this product do not use more than directed</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ nervousness, dizziness, or sleeplessness occur ■ symptoms do not get better within 7 days or occur with fever ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than 6 doses in any 24-hour period ■ measure only with dosing cup provided ■ do not use dosing cup with other products ■ dose as follows or as directed by a doctor ■ adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours ■ children under 12 years of age: do not use <p>Other information</p> <ul style="list-style-type: none"> ■ each 20 mL contains: sodium 12 mg ■ store at 20-25°C (68-77°F) ■ do not refrigerate <p>Inactive ingredients anhydrous citric acid, edetate disodium, FD&C red no. 40, flavors, glycerin (soy), propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate*, xanthan gum *may contain this ingredient</p> <p>Questions? 1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.</p>
Active ingredients (in each 20 mL)	Purposes					
Dextromethorphan HBr 20 mg...Cough suppressant Guaifenesin 400 mg.....Expectorant Phenylephrine HCl 10 mg.....Nasal decongestant						




Image 2: Detailed Drug Facts label, including warnings and directions.

3. ACTIVE INGREDIENTS

Each 20 ml dose contains:

- **Dextromethorphan HBr (20 mg):** Cough Suppressant
- **Guaifenesin (400 mg):** Expectorant
- **Phenylephrine HCl (10 mg):** Nasal Decongestant

3 POWERFUL ACTIVE INGREDIENTS FOR MULTI-SYMP TOM RELIEF

- ▶ **DEXTROMETHORPHAN HBR**
- ▶ **GUAIFENESIN**
- ▶ **PHENYLEPHRINE HCl**

Image 3: Mucinex Fast-Max bottle with callouts for the three active ingredients.

4. DIRECTIONS FOR USE

Dosage Instructions:

- Do not take more than 6 doses in any 24-hour period.
- Measure only with the dosing cup provided. Do not use the dosing cup with other products.
- **Adults and Children 12 Years and Older:** Take 20 ml in the dosing cup provided every 4 hours.
- **Children Under 12 Years of Age:** Do not use.



Image 4: Top view of the bottle cap illustrating how to open the child-resistant cap.

5. STORAGE AND HANDLING

Store at controlled room temperature, between 20-25°C (68-77°F). Do not refrigerate. Keep the bottle tightly closed when not in use.

6. PRODUCT SPECIFICATIONS

Brand	Mucinex
Item Model Number	6380000000
ASIN	B0057UUGWA
Product Dimensions	1.44 x 2.51 x 5.77 inches
Item Weight	0.51 Pounds
Volume	6 Fluid Ounces (180 ml)
Manufacturer	Reckitt Benckiser Healthcare
Date First Available	June 23, 2011



**PEEL CORNER TO READ COMPLETE
DRUG FACTS AND INFORMATION**

Tamper evident: Do not use if neckband
on bottle cap is broken or missing.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



3 63824 01465 4

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224
©2024 RB Health
110624

LOT:
3318170

EXP:

MADE IN:

Image 5: Back of the Mucinex Fast-Max bottle, showing the barcode and manufacturing information.

7. SUPPORT AND CONTACT INFORMATION

For questions or further assistance, please contact Mucinex directly.

Customer Service: 1-866-MUCINEX (1-866-682-4639)

For medical concerns, always consult your healthcare provider.