Saved from manuals.plus.
Open the live page for links to other documents and user manuals.

Mucinex ASA-100

Mucinex Fast-Max Day & Night Cold & Flu Liquid Medicine Instruction Manual

Model: ASA-100

Product Overview

Mucinex Fast-Max Day Time Cold & Flu and Night Time Cold & Flu Liquid Medicine is formulated to provide multi-symptom relief for cold and flu symptoms. This product includes two distinct liquid formulations: one for daytime use and one for nighttime use.
The Day Time formula addresses cough, chest congestion, nasal and sinus congestion, sinus pressure, fever, body pain, headache, and sore throat. The Night Time formula addresses cough, nasal congestion, fever, body pain, headache, sneezing, runny nose, and sore and itchy throat, also aiding in sleep.

Active Ingredients

Day Time Cold & Flu (per 20 mL)

- **Acetaminophen (650 mg):** Pain reliever and fever reducer.
- **Phenylephrine HCl (10 mg):** Nasal decongestant.
- **Dextromethorphan HBr (20 mg):** Cough suppressant.
- **Guaifenesin (400 mg):** Expectorant, helps thin and loosen mucus.

Night Time Cold & Flu (per 20 mL)

- **Acetaminophen (650 mg):** Pain reliever and fever reducer.
- **Phenylephrine HCl (10 mg):** Nasal decongestant.
- **Diphenhydramine HCl (25 mg):** Cough suppressant and antihistamine.

Dosage and Administration

Always use the provided dosing cup for accurate measurement. Do not use other measuring devices.

- **Adults and children 12 years of age and older:** Take 20 mL every 4 hours.
- Do not take more than a total of 6 doses in any 24-hour period.
- Do not take the first dose of Mucinex Fast-Max Night Time Cold & Flu liquid sooner than 4 hours after the last dose of

Mucinex Fast-Max Day Time Severe Cold liquid, unless directed by a doctor.

- **Children under 12 years of age:** Do not use this product.

Each 20 mL dose contains 12 mg of sodium.

Warnings and Precautions

It is important to read all warnings and directions before use. Consult a doctor or pharmacist if you have any questions or pre-existing medical conditions.

- **Overdose Warning:** Taking more than the recommended dose can cause serious liver damage. In case of overdose, seek medical help or contact a Poison Control Center right away.
- **Alcohol Warning:** Avoid alcoholic drinks while taking this product.
- **Do not use** with any other drug containing acetaminophen (pain reliever/fever reducer), as this can lead to overdose.
- **Do not use** if you are allergic to any of the ingredients.
- **Ask a doctor before use if you have:** liver disease, heart disease, high blood pressure, thyroid disease, diabetes, glaucoma, difficulty in urination due to enlarged prostate gland, chronic cough, cough that occurs with too much phlegm (mucus), or asthma.
- **Ask a doctor or pharmacist before use if you are:** taking blood thinning drug warfarin, or taking sedatives or tranquilizers (Night Time only).
- **Stop use and ask a doctor if:** pain or cough gets worse or lasts more than 7 days, fever gets worse or lasts more than 3 days, redness or swelling is present, new symptoms occur, or nervousness, dizziness, or sleeplessness occur (Night Time only).
- **Pregnancy and Breast-feeding:** If pregnant or breast-feeding, ask a health professional before use.
- **Keep out of reach of children.**
- **Driving/Operating Machinery:** The Night Time formula may cause drowsiness. Exercise caution when driving a motor vehicle or operating machinery.

Storage Instructions

Store the product between 20-25°C (68-77°F). Do not refrigerate.

Product Visuals



Image: Front view of the Mucinex Fast-Max Day Time Cold & Flu and Night Time Cold & Flu liquid medicine bottles in their combined packaging.

Drug Facts

Active ingredients (in each 20 mL) Purposes
Mucinex Fast-Max DAY TIME COLD & FLU

Acetaminophen 650 mg	**Pain reliever/fever reducer**
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Active ingredients (in each 20 mL) Purposes
Mucinex Fast-Max NIGHT TIME COLD & FLU

Acetaminophen 650 mg	**Pain reliever/fever reducer**
Diphenhydramine HCl 25 mg	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

DAY TIME COLD & FLU

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - stuffy nose
  - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

NIGHT TIME COLD & FLU

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - sinus congestion and pressure
  - runny nose
  - sneezing
  - itching of the nose or throat
  - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen.
Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on the skin ***(NIGHT TIME only)***
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma ***(NIGHT TIME only)***
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis ***(NIGHT TIME only)***
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers ***(NIGHT TIME only)***

When using this product

- do not use more than directed
- excitability may occur, especially in children ***(NIGHT TIME only)***
- marked drowsiness may occur ***(NIGHT TIME only)***
- alcohol, sedatives, and tranquilizers may increase drowsiness ***(NIGHT TIME only)***
- avoid alcoholic drinks ***(NIGHT TIME only)***
- be careful when driving a motor vehicle or operating machinery ***(NIGHT TIME only)***

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: **sodium 12 mg**
- store between 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients (Mucinex Fast-Max DAY TIME COLD & FLU) anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate†, xanthan gum
†may contain this ingredient

Inactive ingredients (Mucinex Fast-Max NIGHT TIME COLD & FLU) anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate‡, xanthan gum
‡may contain this ingredient



DOSAGE INFORMATION

ADULTS AND CHILDREN 12 YEARS AND OVER:

Take 20 mL in dosing cup provided every 4 hours.

Do not take more than 6 doses in any 24-hour period.

CHILDREN UNDER 12 YEARS:

Do not use

Use as directed.

Image: Detailed view of the Mucinex Fast-Max Day and Night packaging, highlighting dosage information for adults and children 12 years and older.



Image: Close-up of the Drug Facts panel on the Mucinex Fast-Max packaging, showing active ingredients, purposes, warnings, and directions.



Image: Chart illustrating the symptoms relieved by the Day Time and Night Time formulas, including active ingredients for each.



Image: Graphic indicating that one dose relieves 9 symptoms, with visual representations for Day and Night relief.

Product Videos

Mucinex FastMax All in One for Cold & Flu

▶ **Watch video online**

Mucinex FastMax All in One for Cold & Flu

Video: An overview of Mucinex Fast-Max All in One for Cold & Flu, demonstrating its multi-symptom relief capabilities.

Mucinex All In One DJ

▶ **Watch video online**

Mucinex All In One DJ

Video: A short clip featuring the Mucinex All In One product in a dynamic setting, illustrating its effectiveness in helping users return to

their daily activities.

Mucinex AIO Animation only 6s

▶ **Watch video online**
Mucinex AIO Animation only 6s

Video: A brief animated sequence showcasing the Mucinex All In One product.

Specifications

Item Model Number	ASA-100
Product Dimensions	1.44 x 5.01 x 5.77 inches
Item Weight	12 Ounces
Date First Available	August 12, 2013

Manufacturer Information

Manufacturer: Reckitt Benckiser

Legal Disclaimer

Statements regarding dietary supplements have not been evaluated by the FDA and are not intended to diagnose, treat, cure, or prevent any disease or health condition.

For further assistance, please refer to the product packaging or consult a healthcare professional.

Ask a question about this manual

Ask about setup, troubleshooting, compatibility, parts, safety, or missing instructions. Manuals+ will review the question and use this page's manual context to help answer it.

Name

Anonymous

Email

you@example.com

After a successful submission, we securely hash the supplied account or form email before sending it to Microsoft Advertising for conversion attribution. Privacy details.

Question

Example: How do I reset this device or fix this error?

Details

Model number, symptoms, what you tried, or the section of the manual you are using.

Submit question

Open the live page

https://manuals.plus/asin/B00EFRV7Y2

More documents and user manuals: **https://manuals.plus/**