

## Manuals+

[Q & A](#) | [Deep Search](#) | [Upload](#)

manuals.plus /

› [Mucinex](#) /

› [Mucinex Children's Multi-Symptom Cold Liquid: Very Berry Flavor Instruction Manual](#)

## Mucinex B01LY0ILFD

# Mucinex Children's Multi-Symptom Cold Liquid: Very Berry Flavor Instruction Manual

Brand: Mucinex | Model: B01LY0ILFD

## 1. OVERVIEW

Mucinex Children's Multi-Symptom Cold Liquid, Very Berry Flavor, is formulated to help maintain comfort during seasonal changes. This product is available in a convenient multi-pack containing four 4 fl oz (118 mL) bottles, offering both Day Time and Night Time formulas. The liquid features a very berry flavor to facilitate administration to children.



Image: Mucinex Children's Multi-Symptom Cold Liquid Value Pack, featuring both Day Time and Night Time formulas in Very Berry flavor. The packaging highlights key benefits and age recommendations.

## Key Benefits:

- **Child-Friendly Flavor:** Very berry flavor designed to be palatable for children.
- **Convenient Multi-Pack:** Includes four bottles for extended availability.
- **Easy-to-Pour Bottle:** Designed with a child-safe cap and easy-pour format.

## 2. ACTIVE INGREDIENTS

### Day Time Multi-Symptom Cold (in each 5 mL):

- Dextromethorphan HBr 5 mg (Cough Suppressant)
- Guaifenesin 100 mg (Expectorant)
- Phenylephrine HCl 2.5 mg (Nasal Decongestant)

### Night Time Multi-Symptom Cold (in each 10 mL):

- Acetaminophen 325 mg (Pain Reliever/Fever Reducer)
- Diphenhydramine HCl 12.5 mg (Antihistamine/Cough Suppressant)
- Phenylephrine HCl 5 mg (Nasal Decongestant)

## 3. USES

### Day Time Multi-Symptom Cold:

- Relieves stuffy nose
- Controls cough
- Relieves chest congestion
- Breaks up mucus

### Night Time Multi-Symptom Cold:

- Relieves stuffy nose
- Controls cough
- Relieves runny nose & sneezing
- Relieves fever & sore throat

## 4. DOSAGE AND ADMINISTRATION

Always shake well before using. Measure only with the dosing cup provided. Do not use the dosing cup with other products. Dose as follows or as directed by a doctor. Do not give more than 6 doses in any 24-hour period.

### Recommended Dosage

Age	Dose (Day Time)	Dose (Night Time)	Frequency
Children 6 to under 12 years of age	5 mL	10 mL	Every 4 hours
Children 4 to under 6 years of age	2.5 mL	5 mL	Every 4 hours

Age	Dose (Day Time)	Dose (Night Time)	Frequency
Children under 4 years of age	Do not use		

## 5. WARNINGS

- **Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes: more than 5 doses in 24 hours (which is the maximum daily amount) or with other drugs containing acetaminophen.
- **Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include skin reddening, blisters, or rash. If a skin reaction occurs, stop use and seek medical help right away.
- **Sore Throat Warning:** If sore throat is severe, persists for more than 2 days, or is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
- **Do Not Use:**
  - With any other product containing diphenhydramine.
  - With any other product containing acetaminophen.
  - If you are not sure whether a product contains acetaminophen, ask a doctor or pharmacist.
  - If your child is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- **Ask a Doctor Before Use If the Child Has:**
  - Liver disease, heart disease, high blood pressure, thyroid disease, diabetes, glaucoma.
  - A breathing problem such as chronic bronchitis, persistent or chronic cough such as occurs with asthma.
  - Cough that occurs with too much phlegm (mucus).
- **Ask a Doctor or Pharmacist Before Use If the Child Is:**
  - Taking sedatives or tranquilizers.
- **When Using This Product:**
  - Do not use more than directed.
  - May cause excitability, especially in children.
  - Marked drowsiness may occur.
  - Sedatives and tranquilizers may increase drowsiness.
- **Stop Use and Ask a Doctor If:**
  - Nervousness, dizziness, or sleeplessness occur.
  - Symptoms do not get better within 7 days or occur with fever.
  - Cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.
- **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## 6. STORAGE

- Store at 20-25°C (68-77°F).
- Do not refrigerate.

## 7. INACTIVE INGREDIENTS

---

The inactive ingredients include:

- Anhydrous Citric Acid
- Edetate Disodium
- FD&C Blue No. 1
- FD&C Red No. 40
- Flavors
- Glycerin (soy)
- Propyl Gallate
- Propylene Glycol
- Purified Water
- Sodium Benzoate
- Sorbitol
- Sucralose
- Trisodium Citrate Dihydrate
- Xanthan Gum

## 8. TROUBLESHOOTING

---

If symptoms persist or worsen, or if new symptoms appear, consult a healthcare professional. Refer to the 'Warnings' section for specific guidance on when to stop use and seek medical advice.

## 9. SPECIFICATIONS

---

- **Product Name:** Mucinex Children's Multi-Symptom Cold Liquid
- **Flavor:** Very Berry
- **Volume:** 4 Fl Oz (118 mL) per bottle
- **Packaging:** Pack of 4 bottles (Day Time and Night Time formulas)
- **Item Form:** Liquid
- **Age Range (Description):** Kid (specific dosages for 4+ and 6+ years)
- **Manufacturer:** Mucinex
- **ASIN:** B01LY0ILFD

## 10. WARRANTY AND SUPPORT

---

Specific warranty information for this product is not provided in the available data. For any questions or to report side effects, please contact Mucinex customer support.

**Customer Support:** 1-866-MUCINEX (1-866-682-4639)

