

Manuals+

[Q & A](#) | [Deep Search](#) | [Upload](#)

manuals.plus /

› [Scotts](#) /

› [Scotts WeedEx Prevent with Halts: Pre-Emergent Crabgrass and Lawn Weed Preventer Manual](#)

Scotts 10.06 lb. Bag

Scotts WeedEx Prevent with Halts: Pre-Emergent Crabgrass and Lawn Weed Preventer Manual

Model: 10.06 lb. Bag (Covers 5,000 sq. ft.)

1. PRODUCT OVERVIEW



Image: Scotts WeedEx Prevent with Halts product bag on a green lawn.

Scotts WeedEx Prevent with Halts is a granular pre-emergent product designed to prevent the germination of various lawn weeds. This product helps maintain a healthy, weed-free lawn by stopping weeds before they start.


Key Features:

- **Weed Prevention:** Prevents the germination of listed lawn weeds.
- **Season-Long Control:** Effective against crabgrass, foxtail, spurge, barnyardgrass, fall panicum, goosegrass, and oxalis when applied in early spring.
- **Fall Application Benefits:** Prevents moss (except in California), Poa annua, henbit, corn speedwell, and chickweed when applied in the fall.
- **Versatile Use:** Suitable for use on all lawn and grass types, with the exception of Dichondra or Bentgrass lawns.
- **Weather Resistant:** Rain, snow, or freezing weather after application will not affect product performance.
- **Coverage:** One 10.06 lb. bag covers up to 5,000 sq. ft.

2. UNDERSTANDING PRE-EMERGENT WEED CONTROL

Prevents and Controls Problem Weeds*

Including:



Crabgrass



Foxtail



Spurge



Chickweed



Poa Annua

*See label for complete list of weeds

Image: Common weeds prevented by Scotts WeedEx Prevent with Halts.

A pre-emergent weed killer works by forming a barrier in the soil that prevents weed seeds from germinating. It does not kill existing, mature weeds. For best results, it must be applied before weed seeds begin to sprout.

3. WHEN TO APPLY

WHEN TO APPLY

Early spring to prevent crabgrass
and other problem weeds.

During fall to prevent winter weeds
including, Henbit, Chickweed, Poa
Annua, and Corn Speedwell.

Image: Visual guide for application timing.

- **Early Spring:** Apply when the lawn is dry to prevent crabgrass, foxtail, spurge, barnyardgrass, fall panicum, goosegrass, and oxalis. For goosegrass control, a repeat application in 8 weeks may be necessary.
- **Fall (August-October):** Apply to prevent moss (except in California), Poa annua, henbit, corn speedwell, and chickweed.

4. APPLICATION INSTRUCTIONS (SETUP & OPERATING)

Preparation:

- Ensure your lawn is dry before application.
- Identify your lawn size to determine the correct amount of product needed.
- Consult the product bag for specific spreader settings for your Scotts spreader model.

Spreader Settings:

SCOTT'S® WEEDX™ PREVENT WITH HALTS®

provides a simple, easy to use answer for the toughest lawn problems. Their effective ingredients treat the lawn and control the problem. Scott's® WeedEx™ Prevent with Halts® can be used alone, or with other products to solve your lawn problems.

WHAT DOES SCOTT'S® WEEDX™ PREVENT WITH HALTS® DO?

Apply in Spring to prevent crabgrass and other problem weeds. Fall application prevents winter weeds such as: Henbit, Chickweed, Poa Annua and Corn Speedwell.

DIRECTIONS FOR USE | MODO DE EMPLEO

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Se considera una violación a la ley federal usar este producto con otros fines que los indicados en la etiqueta.

Application:

- In early spring, apply when the lawn is dry to prevent germination of crabgrass, foxtail, spurge, barnyardgrass, fall panicum, goosegrass*, and oxalis (common yellow woodsorrel).
- For goosegrass control, repeat application in 8 weeks.
- Apply before March 1 in southern California. For best results on thin lawns apply by mid-February.
- Apply before mid-March in northern California.
- Apply in early Spring (up to time dandelions are in putball stage) in all other states.
- Apply in the fall (August-October) to prevent moss (except in California), or to prevent germination of Poa annua, henbit, corn speedwell, and chickweed.
- For use on all lawns and grass types except dichondra or bentgrass lawns.
- It is not necessary to water-in immediately after application. For best results, lightly water the lawn to rinse particles from grass blades (up to 1/4 to 1/2 inch) within 2-3 days after application.
- Rain, snow or freezing weather after application will not affect the performance of this product.
- This product controls crabgrass in its early stages of growth. This product does not control mature crabgrass and other existing weeds. Growing broadleaf weeds can be killed with Scott's® Turf Builder® Weed & Feed.

This product applies 1.5 lbs of active ingredient per acre. Maximum application rate of 2.0 lbs active ingredient per acre. Can este producto se aplican 0.7 kg de ingrediente activo por acre. La dosis de aplicación máxima es de 0.9 kg de ingrediente activo por acre.

Staining Note: Can cause staining to cement, stone, metallic surfaces (such as automobiles and garden furniture), clothing or shoes. If contact occurs, brush off particles to prevent staining and/or unwanted tracking into the house. If necessary, rinse off with water. Advertencia sobre manchas: Este producto puede causar manchas en superficies de cemento, piedra o metal (tales como automóviles y muebles de jardín), la ropa, o los zapatos. De entrar en contacto con el producto, cepille las partículas para evitar dejar manchas o marcas indeseables al entrar a la casa. De ser necesario, enjuague con agua.

Aplicación:

- A principios de primavera, aplíquelo cuando el césped está seco para prevenir la germinación de digitaria o zacate mano de carraño, zacate cola de zorra, lirrago, capón amaro, pasto colchón, pie de gallina* y vinagrillo (oxalis común amarilla).
- Para controlar el pie de gallina, repita la aplicación o los 8 semanas.
- Aplíquelo antes del 1 de marzo en el sur de California. Para obtener los mejores resultados en céspedes raras, aplíquelo por mediados de febrero.
- Aplíquelo antes de mediados de marzo en el norte de California.
- Aplíquelo a principios de primavera hasta el momento en que los dientes de León estén en la etapa de boriá en todos los demás estados.
- Aplíquelo en el otoño (entre agosto y octubre) para prevenir el musgo (excepto en California) o la germinación de pasto anual, ortiga mansa, verónica del maíz y estalaria.
- Para su uso en todos tipos de céspedes y gramas, excepto oreja de ratón o agrosil.
- No es necesario regar inmediatamente después de la aplicación. Para obtener los mejores resultados, riegue ligeramente para enjuagar las partículas de las bridas del césped (de hasta 0.6 a 1.3 cm) 2 a 3 días después de la aplicación.
- La lluvia, la nieve ni las temperaturas bajo cero afectarán el rendimiento de este producto después de su aplicación.
- Este producto controla el zacate mano de carraño en sus primeras etapas de crecimiento. Este producto no controla el zacate mano de carraño maduro ni otras malezas ya presentes. Las malezas de hoja ancha pueden eliminarse con el Herbicida y fertilizante, para césped Turf Builder® de Scott's®.

1 | SPREADER SETTINGS POSICIONES DEL ESPARCIDOR

SCOTT'S® BROADCAST/ROTARY SPREADER



2 1/4

SCOTT'S® DROP SPREADER



4 1/2

SCOTT'S® WIZZ® SPREADER



3 1/4

2 | COVERAGE TIPS CONSEJOS DE COBERTURA

BROADCAST/ROTARY SPREADER & WIZZ® SPREADER

Esparcidor giratorio/por voleo y esparcidor Wizz®



Make each pass about 5 feet from previous pass. Realice cada pasada a aproximadamente 1.5 metros de distancia de la anterior.

DROP SPREADER

Esparcidor por gravedad



Overlap wheel marks to ensure uniform coverage. Traslape las huellas dejadas por la rueda para asegurar una cobertura uniforme.

RESTRICTIONS | RESTRICCIONES

Do not enter or allow others to enter the treated area until dust has settled. Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. DON'T use on newly seeded grass lawns until after the fourth mowing. On newly seeded areas, wait 5 months before application. DON'T plant grass seed for four growing months or spring an area for 5 months after use of this product. DON'T apply more frequently than twice a year. Allow at least a two month interval between applications. DON'T rake or aerify after application, as the Scott's® WeedEx™ Prevent with Halts® chemical crabgrass barrier in the soil will be disturbed.

No entre ni permita que nadie entre al área tratada hasta que el polvo se haya asentado. No aplique este producto de forma que entre en contacto con otras personas o mascotas, ya sea directamente o al ser arrastrado por el viento. Durante la aplicación, mantenga alejados del área a las personas y las mascotas. NO utilice en prados recién sembrados sino hasta después del cuarto corte de césped. En áreas recién plantadas, espere 5 meses antes de la aplicación. NO sembrar semillas de césped durante cuatro de los meses de crecimiento ni plantar en el área durante cinco meses después de utilizar este producto. NO aplicar con mayor frecuencia que dos veces por año. Deje pasar un intervalo de por lo menos dos meses entre cada aplicación. NO rabillo ni aire la tierra después de la aplicación, pues esto trastocará la barrera química de Scott's® WeedEx™ Prevent with Halts® contra el pasto mano de carraño en la tierra.

SCOTT'S.COM

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION Causes eye irritation. Avoid contact with eyes, skin or clothing. May be harmful if swallowed. Avoid inhalation.

USER SAFETY RECOMMENDATIONS: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.
If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS: This pesticide is toxic to fish. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweep any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems. In case of spills, either collect for use or dispose of properly.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a dry place. Re-seal opened bag by folding top down and securing.
PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.
If empty: Place in trash or other for recycling if available.
If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE To the extent consistent with applicable law, buyer assumes all risk of use, storage or handling of this product not in accordance with directions.

AVISO En la medida contemplada por las leyes correspondientes, el comprador asume todos los riesgos de uso, almacenamiento o manejo de este producto si no se siguen las instrucciones.



If for any reason you, the consumer, are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

DEDICATED TO A BEAUTIFUL WORLD™
Trusted by generations of homeowners, Scott's® is dedicated to helping you enhance and protect the beauty of your home environment.

AVISOS DE PRECAUCIÓN

RIESGOS PARA LOS SERES HUMANOS Y LOS ANIMALES DOMÉSTICOS

PRECAUCIÓN Causa irritación en los ojos. Evite el contacto con los ojos, la piel o la ropa. Puede ser nocivo si se ingiere. Evite inhalarlo.

RECOMENDACIONES DE SEGURIDAD PARA EL USUARIO: Lavarse las manos antes de comer, beber, masticar chicle, consumir tabaco o usar el baño. Los usuarios deberán quitarse la ropa de inmediato si les cae dentro pesticida, para luego lavarse bien y ponerse ropa limpia.

PRIMEROS AUXILIOS

Si entra en los ojos: Mantenga el ojo abierto y enjuáguelo lentamente y con cuidado con agua por 15 a 20 minutos. Retire las lentes de contacto, de haberlas, después de transcurridos los primeros 5 minutos, y continúe enjuagando el ojo.
En caso de contacto con la piel o la ropa: Quitar la ropa contaminada. Enjuague la piel de inmediato con abundante agua por 15 a 20 minutos.
Si se ingiere: Llame a un centro de control de envenenamientos o a un médico de inmediato para recibir instrucciones de tratamiento. Que la persona beba un vaso de agua si puede tragar. No induzca el vómito, a menos que así se lo indique el centro de control de envenenamientos o un médico. No le dé nada por la boca a una persona que haya perdido el conocimiento.
Si se inhala: Lleve a la persona al aire fresco. Si la persona no respira, llame al 911 o a una ambulancia y luego, de ser posible, déle respiración artificial (boca a boca, preferiblemente). Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento. Tenga el envase o la etiqueta del producto con usted cuando llame al centro de control de envenenamientos o al médico, o al acudir a recibir tratamiento.

RIESGOS AMBIENTALES: Este pesticida es tóxico para los peces. Los escurremientos desde las áreas tratadas pueden ser peligrosos para los organismos acuáticos de las zonas adyacentes. Para proteger el medio ambiente, no permita que el pesticida penetre o se escurra en drenajes pluviales, zanjas de drenaje, cunetas o aguas superficiales. La aplicación de este producto en un clima tranquilo, cuando no se pronostican lluvias para las próximas 24 horas, asegurará que ni el viento ni la lluvia arrastren o laven el pesticida fuera del área de tratamiento. Barrer cualquier producto que caiga sobre la entrada del automóvil, la acera o la roña de regreso al área de tratamiento, así como a cuerpos de agua o sistemas de drenaje. En caso de derrames, recoja el producto para usarlo o deséchelo como se deba.

ALMACENAMIENTO Y ELIMINACIÓN

ALMACENAMIENTO DEL PESTICIDA: Guardar en un lugar seco. Cierre la bolsa abierta doblando la parte superior y asegurándola.
ELIMINACIÓN DEL PESTICIDA Y MANEJO DEL ENVASE: Este envase no se puede reutilizar. No vuelva a usar ni a llenar este envase.
Si está vacío: Tirarlo a la basura o, si está disponible, ofrécizalo para reciclaje.
Si está parcialmente lleno: Llame a su agencia local de desechos sólidos para obtener instrucciones sobre cómo eliminarlo. Nunca elimine el producto no utilizado en drenajes internos o externos.

SMG Sold By/Comercializado por:
The Scotts Company
14111 Scottslawn Rd.
Marysville, OH 43041

©2018 The Scotts Company LLC
World rights reserved.
Derechos mundiales reservados.

QUESTIONS/INFORMATION Preguntas/Información

Consumer Helpline
Línea de asistencia al consumidor
1-800-543-TURF (8873)
Visit our website/Visite nuestro sitio web
www.scotts.com

EPA Reg. No. 538-192
EPA Est. No. 538-041-2
85053-04-001, 538-04-2
82757-04-001, 9199-04-1
081323-04-001, 82757-FL-001,
084179-7X-001
Superscript used in first letter of run code on end of container.

Image: Detailed spreader settings from the product bag.

Always refer to the back of your Scotts WeedEx Prevent with Halts bag for the most accurate and up-to-date spreader settings for your specific Scotts spreader model.

Application Steps:



Easy to Use

1. Use on any lawn size

1 bag treats up to
5,000 sq. ft.

2. Fill and set spreader

See label for spreader
settings.

3. Apply to a DRY lawn

Walk at a steady pace.

4. Love your lawn

See product label for
complete instructions.

Image: Easy-to-follow application steps.

1. **Determine Lawn Size:** One 10.06 lb. bag treats up to 5,000 sq. ft.
2. **Fill Spreader:** Pour the granules into your Scotts spreader.
3. **Set Dial:** Adjust your spreader's dial to the recommended setting found on the product bag.
4. **Apply to a DRY Lawn:** Walk at a steady pace to ensure even coverage.
5. **Sweep Hard Surfaces:** If any product lands on cement, stone, or metallic surfaces, sweep it back into the treated lawn area immediately to prevent staining.
6. **Water After Application:** For best results, the lawn should receive 1/4 to 1/2 inch of water from rainfall or light watering within 2-3 days after application.



**USE WITH
A SCOTTS®
SPREADER**

**1 bag treats up to
5,000 SQ. FT.**



Image: Different Scotts spreader types for various lawn sizes.

Application Video:

Your browser does not support the video tag.

Video: Official Scotts guide on how to apply WeedEx Prevent with Halts.

5. COVERAGE

One 10.06 lb. bag of Scotts WeedEx Prevent with Halts is formulated to cover an area of up to 5,000 square feet. Accurately measuring your lawn size before application ensures proper coverage and effectiveness.

6. IMPORTANT CONSIDERATIONS & SAFETY

Weather Conditions:



WORRY-FREE WEATHER PROTECTION

Rain, snow or freezing weather
after application will not affect the
performance of this product.

Image: Product is effective even after rain, snow, or freezing weather.

The performance of Scotts WeedEx Prevent with Halts will not be affected by rain, snow, or freezing weather occurring after application.

Staining Prevention:

- This product could cause staining on cement, stone, or metallic surfaces, as well as clothing and shoes.
- To avoid stains, sweep any stray particles off hard surfaces and back into the treated lawn area immediately.
- Brush off shoes and clothing thoroughly after application. If necessary, use water to rinse particles away from hard surfaces.

Grass Types:

Scotts WeedEx Prevent with Halts is safe for use on all common lawn and grass types, with the specific exceptions of Dichondra or Bentgrass lawns.

Product Limitations:

This product is a pre-emergent weed and crabgrass killer. It controls crabgrass in its early stages of growth by preventing germination. It is important to note that it does not control mature crabgrass or other existing weeds

that have already sprouted.

7. SPECIFICATIONS

Attribute	Value
Brand	Scotts
Item Weight	10.3 Pounds
Item Form	Bag
Active Ingredients	Pendimethalin
UPC	032247490245
Manufacturer	The Scotts Company LLC
Unit Count	160.0 Ounce
Coverage	5000 square_feet

8. WARRANTY AND SUPPORT

For specific warranty information, product guarantees, or further assistance, please refer to the product packaging or visit the official Scotts website. You can also contact Scotts customer service directly for personalized support regarding application, product performance, or any other inquiries.

Scotts Website: www.scotts.com

Customer Service: Refer to product packaging for phone numbers or contact forms.