Saved from manuals.plus.
Open the live page for links to other documents and user manuals.

Mucinex 3135268

Mucinex Fast-Max Cold & Flu and Nightshift Severe Cold & Flu Adult Medication Manual

Model: 3135268 | Brand: Mucinex

Product Overview

The Mucinex Fast-Max Cold & Flu and Mucinex Nightshift Severe Cold & Flu combo pack provides multi-symptom relief for cold and flu symptoms. The daytime formula addresses fever, sore throat, headache, body pain, cough, chest congestion, nasal congestion, and sinus pressure. The nighttime formula helps reduce fever and relieves sore throat, headache, body pain, cough, nasal congestion, sneezing, and runny nose for nighttime symptom management. This product is intended for adults and children 12 years of age and over.



Image: Mucinex Fast-Max Cold & Flu (white bottle) and Mucinex Nightshift Severe Cold & Flu (blue bottle) packaging.

▶ **Watch video online**

Mucinex FastMax All in One for Cold & Flu

Video: An official Mucinex video explaining the benefits of Mucinex Fast-Max All in One for Cold & Flu.

Important Safety Information

Always read and follow all directions and warnings on the product packaging before use. Consult a healthcare professional if you have any questions or concerns.

Keep Out of Reach of Children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Overdose Warning

Taking more than the recommended dose (overdose) may cause liver damage.

Do Not Use If:

- You are taking any other drug containing acetaminophen (prescription or nonprescription).
- You are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- You are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a Doctor Before Use If You Have:

- Liver disease
- Heart disease
- High blood pressure
- Thyroid disease
- Diabetes
- Glaucoma
- Trouble urinating due to an enlarged prostate gland
- Breathing problem such as emphysema or chronic bronchitis
- Persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- Cough that occurs with too much phlegm (mucus)

Ask a Doctor or Pharmacist Before Use If You Are:

- Taking the blood thinning drug warfarin
- Taking sedatives or tranquilizers

Stop Use and Ask a Doctor If:

- Nervousness, dizziness, or sleeplessness occur
- Pain, nasal congestion, or cough gets worse or lasts more than 7 days
- Fever gets worse or lasts more than 3 days
- Redness or swelling is present
- New symptoms occur
- Cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

Active Ingredients

Daytime Formula (Mucinex Fast-Max Cold & Flu)

Active Ingredient (in each 20 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Nighttime Formula (Mucinex Nightshift Severe Cold & Flu)

Active Ingredient (in each 20 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant
Triprolidine HCl 2.5 mg	Antihistamine



Image: Detailed Drug Facts label for Mucinex Fast-Max Cold & Flu, showing active ingredients and warnings.



NIGHTTIME SEVERE COLD & FLU

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant
Triprolidine HCl 2.5 mg	Antihistamine

Uses

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - sinus congestion and pressure
  - runny nose
  - sneezing
  - itching of the nose or throat
  - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product ■ do not use more than directed

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

Tamper evident: Do not use if neckband on bottle cap is broken or missing.

COLD & FLU

Drug Facts

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
  - cough
  - nasal congestion
  - minor aches and pains
  - sore throat
  - headache
  - stuffy nose
  - sinus congestion and pressure
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 6 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Image: Detailed Drug Facts label for Mucinex Nightshift Severe Cold & Flu, showing active ingredients and warnings.

Directions for Use

Dosage for Adults and Children 12 Years of Age and Over:

- Take 20 mL in dosing cup provided every 4 hours.
- Do not take more than 4 doses in any 24-hour period.

Important Notes:

- Measure only with dosing cup provided.
- Do not use with other products containing the same active ingredients.
- Do not refrigerate.
- Always wait at least 4 hours before taking another dose of Mucinex liquid.



Image: List of symptoms relieved by Mucinex Fast-Max Cold & Flu daytime formula, including headache, body pain, sore throat, cough, sinus pressure, fever, chest congestion, and sinus congestion.



Image: List of symptoms relieved by Mucinex Nightshift Severe Cold & Flu nighttime formula, including headache, body pain, sore throat, nasal congestion, cough, fever, and sneezing.

Inactive Ingredients

Mucinex Fast-Max Cold & Flu: Anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum. May contain this ingredient.

Mucinex Nightshift Severe Cold & Flu: Anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose, trisodium citrate dihydrate, xanthan gum. May contain this ingredient.

Questions and Support

For questions or to report side effects, please call 1-866-MUCINEX (1-866-682-4639).

Visit our website for more information:www.mucinex.com

Patent information: www.rb.com/patents

Product Specifications

- **Product Dimensions:** 1.44 x 5.01 x 5.77 inches
- **Item Weight:** 0.78 Pounds (12.48 ounces)
- **Item Model Number:** 3135268
- **Date First Available:** August 6, 2020
- **Manufacturer:** Reckitt Benckiser
- **Specific Uses For Product:** Cold, Flu
- **Product Benefits:** Cough, Cold & Flu Relief, Sore Throat, Fever, & Congestion Relief, Pain Relief, Sinus Control, Headache Relief

Additional Information



Image: A visual representation of the 'All in One' relief provided by Mucinex, showing various symptoms addressed.



Image: Mr. Mucus character alongside text indicating '12 Total Fl Oz (2 x 6 Fl Oz bottles)'.



Image: Text 'NOTHING WORKS FASTER*' on a blue background, with a small disclaimer '*AMONG ORAL OTC COLD/FLU LIQUIDS'.



Image: A display of various Mucinex products, each targeting specific symptoms like cough, chest congestion, and sinus relief.

COLD & FLU (cont'd)

Drug Facts (continued)

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period



Image: Illustration showing that product packaging may vary, displaying two different designs for the Mucinex Day & Night Cold & Flu pack.



- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- adults and children 12 years of age and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

Other information

- each 20 mL contains: **sodium 16 mg**
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

ammonium glycyrrhizate, anhydrous citric acid, ascorbic acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin (soy), propylene glycol, sodium benzoate, sorbitol, sucralose, triacetin, triethyl citrate, water, xanthan gum

Questions?

1-866-MUCINEX (1-866-682-4639)

POWERFUL RELIEF FOR YOUR NIGHT
+
POWERFUL RELIEF FOR YOUR DAY

reckitt

Dist. by: RB Health (US)
Parsippany, NJ 07054-0224



Image: Continuation of Drug Facts for both Cold & Flu and Nighttime Severe Cold & Flu, including directions and inactive ingredients.

▶ **Watch video online**
Mucinex All In One DJ

Video: An official Mucinex video showcasing the product in a dynamic setting.

▶ **Watch video online**
Mucinex FastMax - TVC_Subtitles_101022

Video: An official Mucinex commercial with subtitles, highlighting the product's effectiveness.

© 2024 Mucinex. All rights reserved.

Ask a question about this manual

Ask about setup, troubleshooting, compatibility, parts, safety, or missing instructions. Manuals+ will review the question and use this page's manual context to help answer it.

Name

Anonymous

Email

you@example.com

After a successful submission, we securely hash the supplied account or form email before sending it to Microsoft Advertising for conversion attribution. Privacy details.

Question

Example: How do I reset this device or fix this error?

Details

Model number, symptoms, what you tried, or the section of the manual you are using.

Submit question

Open the live page

https://manuals.plus/asin/B08836KTVQ

More documents and user manuals: **https://manuals.plus/**