

Patient Blood Management: Improved Decision Making With Real Time Data

Sherri Ozawa, RN

Clinical Director, Institute for Patient Blood Management and Bloodless Medicine and Surgery Englewood Hospital and Medical Center, Englewood, NJ

Director, Society for the Advancement of Blood Management, Houston, Texas

May, 2017

Patient Blood Management: Improved Decision Making with Real Time Data

Sherri Ozawa, RN

Clinical Director

Institute for Patient Blood Management and Bloodless

Medicine and Surgery

Englewood Hospital and Medical Center

Englewood, NJ

Director, Society for the Advancement of Blood

Management

Houston, Texas

May, 2017



Learning Objectives

- Address the drivers for change in transfusion practice
- Discuss the importance of modalities for Blood Conservation
- Explain the limitations of invasive hemoglobin measurements
- Introduce the connection between real time hemoglobin data and clinical decision making
- Describe the role of SpHb Monitoring in a comprehensive Patient Blood Management Program

Why do we need to discuss transfusion?

- 1. Frequency – more than we realize*
- 2. Risks- higher than most believe*
- 3. Costs- large and increasing*
- 4. Behavior - non evidence based*
- 5. Overuse - dangerous*




Clinicians' Perceptions


- Safety of BLOOD - **high**
 - Risk of BLOOD - **low**
- Risk of Anemia - **unknown**
- Risk of Surgical Bleeding – **low**
 - Transfusions **rare**



Blood Transfusion: Who is at risk?




Five Drivers for the Paradigm Shift in Transfusion



BLOOD TRANSFUSION

- AHRQ: Blood transfusions occurred in 1/10 of all hospital stays that had a procedure
- Most commonly performed procedure in US hospitals
- It is a Liquid Organ Transplant ordered by a physician and performed by a nurse




US blood collection & consumption trends¹

Allogenic WB / RBC units²

■ Total units collected (M)

— Units transfused per 1000 population



- Blood transfusion rates hovered around 50 units / 1000 members of population from 2001-2008
- Since then, transfusion rates have declined to 40 units / 1000 population by 2013, and experts estimate it may be as low as 35 units today (~30% decline)
- Collection rates have fallen, following a supply glut in 2008

¹ Data available for survey years only; ² ~450-500 mL of whole blood

SOURCE: DHHS National Blood Collection & Utilization Surveys 1997-2011; AABB Blood Survey 2013

Blood Centers: A system in Crisis

MEDICAL ADVANCES RESULT IN A MAJOR
DECLINE IN NEED FOR DONATED
BLOOD COMMUNITY HOSPITALS
DOWN 47%
YEARS, PI
CENTER II

Blood and red ink: Hospitals perform fewer transfusions and blood banks feel the pain



Why blood banks are shrinking



oneblood
Share your power.

DONOR LOGIN & APPOINTMENTS

Join Our E

HOME

MY ACCOUNT

LOCATIONS

LEARN

TARGET YOUR TYPE

STORIES

REWARDS

Media Inquiries

Tell Us Your Story

Press Releases

Graphic Standards, Logos and Images

BLOOD CENTERS ANNOUNCE INTENT TO ME

(Orlando, Florida, July 25, 2014) – OneBlood, Inc., the not-for-profit blood center serving most of Florida and parts of South Georgia and Alabama, and The Institute for Transfusion Medicine, Inc. (ITxM) the not-for-profit blood center and transfusion



New transfusion guidelines, surgical techniques limit need for blood

THE WALL STREET JOURNAL.

[Home](#) [World](#) [U.S.](#) [Politics](#) [Economy](#) [Business](#) [Tech](#) [Markets](#) [Opinion](#) [Arts](#) **Life** [Re](#)



20 ODD QUESTIONS
Designer Jonathan
Adler on Awesome
Architecture ...



The Price We Pay for
Sitting Too Much



New Devices Take
the Pain Out of
Hospital Visits




New Sup
Home Bi

[LIFE](#) | [HEALTH](#) | [JOURNAL REPORTS: HEALTH CARE](#)

The Rising Risk of a Contaminated Blood Supply

Amid new pathogens and cost pressures, efforts are under way to keep patients safe



Patient blood

Table 1 Reported adverse outcomes associated with transfusion and populations affected

Patient populations in whom adverse outcomes associated with RBC transfusion have been reported

ICU patients

Paediatric patients

Trauma patients

Burn patients

Combat casualties

Acute coronary syndrome patients

Oncology patients

Population based studies

Obstetrics

Cardiac surgery

Neurosurgery

Hepatic surgery

Pancreatic surgery

Transplantation surgery

Colorectal surgery

Gastric surgery

Biliary surgery

Splenectomy

General surgery

Vascular surgery

Orthopaedic surgery

Thoracic aneurysm surgery

Lung surgery

Head and neck surgery

Mastectomy and reconstruction

Oesophageal surgery

Transfusion

Ischemia

Postoperative

Recovery

Requiring re-operation

Admission to ICU


Length of stay

Hospital length of stay

Hospital readmission

Mortality


Activity Based Cost of Transfusion from a Provider's Perspective



ICCTO

- > 30,000 citations – 494 analyzed – 450 clinical scenarios
- Appropriateness of Allogeneic Blood Transfusion based on improving health outcomes

ALL “appropriately” transfused patients had Hb less than 7.9 g/dL
No one appropriate if Hb $\geq$ 8.0 g/dL



Blood use in elective surgery: the Austrian benchmark study





Hans Gombotz, Peter H. Rehak , Aryeh Shander , and Axel Hofmann

Who really gets transfused?


97.4% of all transfusions could have been predicted by:

1. Level of anaemia prior to surgery
2. Volume of perioperative blood loss
3. Transfusion trigger

Morphology of RBC's in Stored Blood



Patient Blood Management



Why do we test for hemoglobin?

- To check overall health
- To diagnose a medical condition
- To monitor a medical condition

<http://www.mayoclinic.org/tests-procedures/hemoglobin-test/basics/why-its-done/prc-20015022>

Why do we test for hemoglobin?

- **It is a predictor of transfusion**
- **Delta Hg is associated with worse outcomes**
- **It has the ability to determine if any intervention is required**
- **Determine that an endpoint has been reached**

Delivery of Oxygen to Tissues

Oxygen Delivery Cardiac Output Oxygen Content

$$DO_2 = CO \times CaO_2$$

SV
Stroke Volume

HR
Heart Rate

Hgb
Hemoglobin Concentration

SaO₂
Oxygen Saturation


Dissolved Oxygen

Hemoglobin concentration is a significant factor of oxygen delivery to the tissues.

Decreased levels of hemoglobin can have detrimental effects in a patient due to limited delivery of oxygen to vital organs.¹

Hemoglobin Levels

- Low hemoglobin values can have multiple causes¹
 - Diseases that can cause the body to produce fewer red blood cells
 - Diseases that cause the body to destroy red blood cells
 - Procedures or conditions that cause blood loss
 - Examples: a bleeding wound, GI bleeds from ulcers, etc.



Normal ranges for women and men¹

- > **For more severe blood loss, red blood cell (RBC) transfusion is a common method to increase hemoglobin count in the body**
 - The decision to transfuse is generally based on the value of hemoglobin as well as the **clinical condition of the patient**²

¹World Health Organization Global Database on Anaemia. 2008.

²Miller's Anesthesia. 8th Edition. 2015.

Different types of Hg measurements

- Lab analysis - Automatic blood analyzer/ CO-Oximeters are the gold standard for Hb measurement
- Capillary blood tube - spun
- POCT – Hemocue - reagent
- Non-invasive Hg monitoring
- Visual color-based POC anemia – pre-investigation

Invasive Hemoglobin Methods



- Cyanmethemoglobin (HiCN)
 - International gold standard - Complex, manual process



- Hematology analyzer
 - Best available measurement in most hospitals and reference labs - Brands: Beckman Coulter, Sysmex, Abbott, Bayer, Siemens
- CO-Oximeter
 - Acceptable measurement, less accurate at higher Hb levels - Brands: include Radiometer, Nova Biomedical




- Point-of-care
 - Least accurate invasive method but benefit is quicker test result - Brands: Hemocue, iStat



Limitations of Traditional Methods

Lab Hb Gives Intermittent and Delayed Results While Transfusion
Decisions Are Often Made In Real Time




¹Frank S et al. Anesthesiology. 2012.

²Shander et al. TransMed Rev. 2011.

Limitations of Current Methods

Lab Hb Gives Intermittent and Delayed Results While Transfusion Decisions
Are Often Made In Real Time



Testing vs. monitoring – What is the difference

- **TESTING:** A medical test is a kind of medical procedure performed to detect, diagnose, or monitor diseases, disease processes, susceptibility, and determine a course of treatment.
- **MONITORING:** In medicine, monitoring is the observation of a disease, condition or one or several medical parameters over time.

[http://en.wikipedia.org/wiki/Monitoring_\(medicine\)](http://en.wikipedia.org/wiki/Monitoring_(medicine))


http://en.wikipedia.org/wiki/Medical_test

Hemoglobin monitoring vs hemoglobin testing

- Hemoglobin testing:
 - Reliable
 - “spot check”
 - Relatively long turnaround time
- Hemoglobin monitoring:
 - Continuous real-time monitoring and trending
 - Provides early decision ability

Value of SpHb Monitoring

Real-time Visibility to Changes-or Lack of Changes- in Hemoglobin between Invasive Blood Sampling




Advantages of non-invasive continuous monitoring

In a study of 20 spine surgery patients (published in Anesthesia-Analgesia), Ron Miller and colleagues noted that non-invasive and continuous Hb monitoring may:

- “Allow immediate and ongoing changing Hb levels to be displayed”
- “Allow a more rapid detection of clinically significant blood loss”
- “Has the potential to significantly improve perioperative transfusion practices”

Value of SpHb Monitoring


Real-time Visibility to Changes-or Lack of Changes- in Hemoglobin
between Invasive Blood Sampling



SpHb Utility

Orthopedic Surgery

% of Patients Receiving RBC Transfusion



- > Randomized trial
- > 327 orthopedic surgery patients
- > Massachusetts General Hospital
- > Found that availability of SpHb monitoring reduced the rate of transfusions when compared to standard care without SpHb
 - 0.6% of the SpHb group received a transfusion
 - 4.5% of the Standard Care group received a transfusion
 - 5.7% of the Retrospective group received a transfusion


Author Conclusion: *“the availability of SpHb decreases inappropriate transfusion (either by preventing an initial transfusion, or the transfusion of additional blood products after a single unit has been delivered).”*

SpHb Utility

Neurosurgery

■ Control Group

■ SpHb Group



- > Prospective cohort study
- > 106 neurosurgical patients
- > Adding SpHb monitoring resulted in decreased blood utilization, while also facilitating earlier transfusions
 - Fewer units of blood transfused
 - Fewer patients receiving more than 3 units
 - A shorter time to transfusion after the need was established

Author Conclusion: “*The ability to observe the continuous trend in hemoglobin affects transfusion behavior, allowing earlier cessation of RBC transfusion as well as earlier consideration of initiation of RBC transfusion.*”

Multi-Level Spine

Utilization of Blood

- About 57% (aggregate) of spinal patients receive blood
 - Ranging from 26% to 79% based upon patient pathology¹
- Clinicians often overestimate the need for blood in spine cases
 - Crossmatch per patient often exceeds amount transfused




Pathology Group	Transfused	Crossmatch/ Patient	Transfuse/ Patient
Deformity (scoliosis, kyphosis)	79%	6	5
Fracture (spinal repair and fusion)	65%	6	4
Tumor (tumor removal)	50%	6	3
Painful Spine (reconstruction and fusion)	26%	3	1
Total	57%	5	3

¹Kouritas et al. Transfusion Med. 2010.

Multi-Level Spine

Case Example- Pediatric Scoliosis



Multi-Level Spine

Peer Reviewed Studies

> Study: **Berkow et al.¹**

- Objective: To evaluate the accuracy of SpHb with tHb during complex spine procedures in patients at high risk for blood loss.
- Sample: 29 patients. Complex spine surgery

> Authors' Conclusions:

- “Continuous, noninvasive hemoglobin measurement via Pulse CO-Oximetry demonstrated clinically acceptable accuracy of hemoglobin measurement within 1.5 g/dL compared with a standard laboratory reference device when used during complex spine surgery.”

> Study: **Colquhoun et al.²**

- Objective: To assess the ability of the Masimo Pulse CO-Oximetry platform to measure changes in Hgb in the intraoperative setting, by comparing SpHb estimates to Hgb from arterial blood gases in major neurosurgical procedures with significant expected blood loss.
- Sample: 20 patients. Major lumbar and low thoracic spine surgery

> Authors' Conclusions:

- “Pulse CO-Oximetry offers an acceptable trend monitor in patients undergoing major spine surgery.”

¹Berkow et al. Anesth, & Analg. 2011.

²Colquhoun et al. J Clin Monit Comput. 2012.

Multi-Level Spine

Peer Reviewed Studies

> **Study: Miller et al.¹**

- Objective: To compare the accuracy of SpHb and HemoCue® with Lab Hemoglobin Values (Coulter Counter).
- Sample: 20 general spine surgery patients

> **Authors' Conclusions:**

- **“We conclude that SpHb could have frequently been used in many patients to guide clinical decisions regarding the need for blood transfusions.”**

¹ Miller et al. Anesthesia & Analgesia. 2011.

HemoCue is a registered trademark of © 2016 HemoCue America a division of Radiometer

Liver Transplants

Utilization of Blood



- Liver Transplants are the bloodiest surgeries
 - Despite low procedure volumes, fall into top 5 for transfusions¹
 - Procedures are long: ranging from 6 to 12 hours²
- 75% of patients who undergo liver surgery are anemic³
 - Liver disease often causes portal hypertension leading to varices and thus gastrointestinal bleeding
 - Endothelial dysfunction impacts clotting abilities and often leads to coagulopathy
 - Alcohol is toxic to bone marrow, thus impacting the ability to regenerate red blood cells
 - Treatments, such as ribavirin-induced hemolysis for Hep-C patients, may cause anemia as a complication
- The risk of bleeding from coagulopathy and iatrogenic injury is high during the early post-transplantation period⁴
 - Roughly 9% (7-15% per Cleveland Clinic) of patients showed abdominal bleeding occurring at a mean of 6.1 days (range 1-21 days)

¹CMS data- ICD 99.0


²https://my.clevelandclinic.org/health/treatments_and_procedures/hic_Liver_Transplantation

³Gonzalez-Casas et al. World J Gastroenterology. 2009.

⁴Jung et al. Transplant Proc. 2012.

Liver Transplant

Case Example



Liver Transplants

Abstracts

> Study: Ramsay et al.¹

- Objective: To evaluate the trend accuracy of SpHb monitoring in a case series of high blood loss, liver transplant surgery patients.
- Sample: 7 patients (30 points). Liver transplants

> Authors' Conclusions:

- **“SpHb monitoring with spectrophotometric sensor provides continuous, precise real-time data on changes in hemoglobin during dynamic conditions such as high blood loss liver transplantation and therefore may be useful in guiding blood management.”**

> Study: Torp et al.²

- Objective: To determine if noninvasive continuous monitoring of tHb using the rainbow SET™ Pulse CO-Oximeter® is an adequate and reliable detector of the trend and magnitude of hemoglobin changes during liver transplant surgery.
- Sample: 15 patients (89 points). Liver transplants

> Authors' Conclusions:

- **“This monitor has the potential to improve the timeliness and precision of transfusion practices and to help clinicians maintain optimal hemoglobin levels.”**

¹Ramsay et al. NYPGA. 2013.

²Torp et al. NYPGA. 2010.

Cranial and Facial

Utilization of Blood

- > **Select cranial and facial procedures (including maxillo) have significant blood loss**
 - Glioma Excision - 834mL blood loss¹
 - CP Angle Tumors - 568mL blood loss¹
 - Decompressive Craniectomy - 338mL blood loss²
 - Craniofacial Surgeries - 410mL blood loss³
 - Double Jaw Osteotomies - 450mL blood loss⁴

- > **Much of the craniofacial data involves pediatric patients⁵⁻⁷**



¹Bhatnagar et al. Indian Journal of Anaesthesia 2007.

²Desgranges et al. Childs Nerv Syst. 2014

³Abraham et al. The Internet Journal of Anesthesiology. 2014.

⁴Moenning et al. J Oral Maxillofac Surg.


⁵Park et al. Anesth, & Analg. 2012.

⁶Patino et al. Anesth & Analg. 2014.

⁷Loveland-Baptist et al. Annual Meeting of the ASA. 2011.

Cranial and Facial

Case Example- Craniectomy



Cranial and Facial

Peer Reviewed Studies

> Study: **Park et al.**¹

- Objective: To compare noninvasive measurements of SpHb with simultaneous laboratory measurements of total hemoglobin in arterial blood samples taken from children (tHb)
- Sample: 40 children (130 points). Brain tumor, craniosynostosis, or moyamoya disease

> Authors' Conclusions:

- **"The Radical-7® Pulse CO-Oximeter can be useful as a trend monitor in children during surgery even immediately after intravascular volume expanders are administered."**

> Study: **Patino et al.**²

- Objective: To assess the trending and accuracy of SpHb with Hb concentration obtained with conventional laboratory techniques (Hb) in children undergoing surgical procedures with potential for substantial blood loss
- Sample: 46 children (158 points). Craniofacial surgery (among others)

> Authors' Conclusions:

- **"SpHb displayed similar trending and absolute accuracy in pediatric patients undergoing a variety of surgical procedures associated with blood loss as has been found in adult surgical patients."**



¹Park et al. Anesth. & Analg. 2012.

²Patino et al. Anesth & Analg. 2014.

SpHb Monitoring Solutions



Root® Monitor



Philips IntelliVue MX800

Conclusion

- Transfusion is complex and costly procedure with clinical and societal implications
- There are multiple drivers for a paradigm shift in transfusion practice
- Traditional Hemoglobin testing has limitations and inefficiencies which can lead to less than optimal clinical decisions
- Non Invasive hemoglobin monitoring (SpHb) provides real time data to clinicians leading to better decision making

