



# prolia Risk Evaluation and Mitigation Strategy Instructions

Home » prolia » prolia Risk Evaluation and Mitigation Strategy Instructions

#### **Contents**

- 1 prolia Risk Evaluation and Mitigation Strategy
- 2 INDICATION
- **3 IMPORTANT SAFETY INFORMATION**
- **4 INDICATION**
- 5 Documents / Resources
  - **5.1 References**
- **6 Related Posts**



# prolia Risk Evaluation and Mitigation Strategy



# **INDICATION**

Prolia® is a prescription medicine used to treat osteoporosis in women after menopause who are at high risk for

fracture or cannot use another osteoporosis medicine or other osteoporosis medicines that did not work well. Here are some key reasons why you may be interested in Prolia®, and important topics to discuss further with your doctor:

Here are some key reasons why you may be interested in Prolia®, and important topics to discuss further with your doctor:

- 1. Only Prolia® is proven to help strengthen and protect bones from fracture with 1 shot every 6 months. You should take calcium and vitamin D as your doctor tells you to while you receive Prolia®. After your treatment with Prolia® is stopped, or if you skip or delay taking a dose, your risk for breaking bones, including bones in your spine, is increased. Do not stop, skip, or delay taking Prolia® without first talking with your doctor.
- 2. Women taking Prolia° for 3 years reduced their risk of new spine fractures by 68%.
  - In a 3-year study, women not treated with Prolia® had more new spine fractures (7.2%) compared to women treated with Prolia® (2.3%). The study consisted of 7393 patients who received either Prolia® or a placebo (a treatment containing no medicine).
- 3. In a 1-year study, women who switched from alendronate to Prolia® showed a significant increase in bone mineral density at the hip and lumbar spine compared to women who stayed on alendronate (a commonly prescribed bisphosphonate, such as Fosamax®).

This study of over 500 women with postmenopausal osteoporosis compared the bone mineral density (BMD) of women who switched from alendronate to Prolia® vs women who stayed on their alendronate treatment. This study only evaluated bone mineral density. These results do not imply fracture risk reduction.



You should take calcium and vitamin D as your doctor tells you to while you receive Prolia®. After your treatment with Prolia® is stopped, or if you skip or delay taking a dose, your risk for breaking bones, including bones in your spine, is increased. Do not stop, skip or delay taking Prolia® without first talking with your doctor. Prolia® is covered by most insurance, including Medicare and many commercial plans. Individual out-of-pocket costs will vary. Talk to your insurance provider to determine what you will pay for Prolia®.

#### **IMPORTANT SAFETY INFORMATION**

#### What is the most important information I should know about Prolia®?

If you receive Prolia®, you should not receive XGEVA® (denosumab). Prolia® contains the same medicine as XGEVA®.

# Prolia® can cause serious side effects (including):

Increased risk of severe low calcium levels in your blood (hypocalcemia). Prolia may lower the calcium levels in your blood. If you have low blood calcium before you start receiving Prolia, it may get worse during treatment. Your low blood calcium must be treated before you receive Prolia. Talk to your doctor before starting Prolia. Your doctor may prescribe calcium and vitamin D to help prevent low calcium levels in your blood while you take Prolia. Take calcium and vitamin D as your doctor tells you to. If you have advanced chronic kidney disease (may or may not be on kidney dialysis), Prolia may increase your risk for severely low calcium levels in your blood, which could result in hospitalization, life-threatening events, and death. A mineral and bone disorder associated with kidney disease called chronic kidney disease-mineral bone disorder (CKD-MBD) may increase your risk for severely low calcium levels in your blood. Before you start PROLIA and during treatment, your doctor may need to do certain blood tests to check for CKD-MBD.

#### Please see additional Important Safety Information on the next page.

2024 Amgen Inc. All rights reserved. USA-162-83526

#### What is the most important information I should know about Prolia®?

If you receive Prolia®, you should not receive XGEVA® (denosumab). Prolia® contains the same medicine as XGEVA®.

#### Prolia® can cause serious side effects (including)

Increased risk of severe low calcium levels in your blood (hypocalcemia). Prolia may lower the calcium levels in your blood. If you have low blood calcium before you start receiving Prolia, it may get worse during treatment. Your low blood calcium must be treated before you receive Prolia. Talk to your doctor before starting Prolia. Your doctor may prescribe calcium and vitamin D to help prevent low calcium levels in your blood while you take Prolia. Take calcium and vitamin D as your doctor tells you to. If you have advanced chronic kidney disease (may or may not be on kidney dialysis), Prolia may increase your risk for severely low calcium levels in your blood, which could result in hospitalization, life-threatening events, and death. A mineral and bone disorder associated with kidney disease called chronic kidney disease-mineral bone disorder (CKD-MBD) may increase your risk for severely low calcium levels in your blood. Before you start PROLIA and during treatment, your doctor may need to do certain blood tests to check for CKD-MBD. Most people with low blood calcium levels do not have symptoms, but some people may have symptoms. Call your doctor right away if you have symptoms of low blood calcium such as:

- spasms, twitches, or cramps in your muscles
- numbness or tingling in your fingers, toes, or around your mouth.

**Serious allergic reactions** have happened in people who take Prolia®. Call your doctor or go to your nearest emergency room right away if you have any symptoms of a serious allergic reaction, including low blood pressure (hypotension); trouble breathing; throat tightness; swelling of your face, lips, or tongue; rash; itching; or hives.

**Unusual thigh bone fractures.** Some people have developed unusual fractures in their thigh bone. Symptoms of a fracture include new or unusual pain in your hip, groin, or thigh.

# Increased risk of broken bones, including broken bones in the spine, after stopping, skipping or delaying Prolia®.

Talk with your doctor before starting Prolia® treatment. After your treatment with Prolia® is stopped, or if you skip or delay taking a dose, your risk for breaking bones, including bones in your spine, is increased. Your risk for having more than 1 broken bone in your spine is increased if you have already had a broken bone in your spine. Do not stop, skip or delay taking Prolia® without first talking with your doctor. If your Prolia® treatment is stopped, talk to your doctor about other medicines that you can take.

**Serious infections** in your skin, lower stomach area (abdomen), bladder, or ear may happen. Inflammation of the inner lining of the heart (endocarditis) due to an infection may also happen more often in people who take Prolia®. You may need to go to the hospital for treatment. Prolia® is a medicine that may affect the ability of your body to fight infections. People who have weakened immune systems or take medicines that affect the immune system may have an increased risk for developing serious infections.

Skin problems such as inflammation of your skin (dermatitis), rash, and eczema have been reported.

Bone, joint, or muscle pain. Some people who take Prolia® develop severe bone, joint, or muscle pain.

**Do not take Prolia®** if you: have low blood calcium; or are pregnant or plan to become pregnant, as Prolia® may harm your unborn baby; or are allergic to denosumab or any ingredients in Prolia®.

Before taking Prolia®, tell your doctor about all of your medical conditions, including if you:

- Take the medicine XGEVA® (denosumab)|
- · Have low blood calcium
- · Cannot take daily calcium and vitamin D
- Had parathyroid or thyroid surgery (glands located in your neck)
- Have been told you have trouble absorbing minerals in your stomach or intestines (malabsorption syndrome)
- · Have kidney problems or are on kidney dialysis
- Are taking medicine that can lower your blood calcium levels
- · Plan to have dental surgery or teeth removed
- · Are pregnant or plan to become pregnant

#### Females who can become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment with Prolia®.
- You should use an effective method of birth control (contraception) during treatment with Prolia® and for at least 5 months after your last dose of Prolia®.
- Tell your doctor right away if you become pregnant while taking Prolia®.
- · Are breast-feeding or plan to breast-feed

#### What are the possible side effects of Prolia®?

It is not known if the use of Prolia® over a long period may cause slow healing of broken bones. The most common side effects of Prolia® are back pain, pain in your arms and legs, high cholesterol, muscle pain, and bladder infection. These are not all the possible side effects of Prolia®. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit <a href="https://www.fda.gov/medwatch">www.fda.gov/medwatch</a>, or call 1-800-FDA-1088.

### **INDICATION**

Prolia® is a prescription medicine used to treat osteoporosis in women after menopause who are at high risk for fracture or cannot use another osteoporosis medicine or other osteoporosis medicines that did not work well. Please see Prolia® full Prescribing Information, including Medication Guide on <a href="mailto:prolia.com">prolia.com</a>.

• 2024 Amgen Inc. All rights reserved. USA-162-83526

#### **Documents / Resources**



<u>prolia Risk Evaluation and Mitigation Strategy</u> [pdf] Instructions
Risk Evaluation and Mitigation Strategy, Evaluation and Mitigation Strategy, Mitigation Strategy,
Strategy

#### References

User Manual